

# Questions to ask about urothelial cancer

## Questions to ask after diagnosis

---

What stage urothelial cancer do I have? Is it non-invasive, invasive or advanced?

Can you explain the results to me?

Are there support services available?

Who will be leading my treatment plan?

## Questions to ask about treatments

---

What treatments are best for me?

What are the goals of these treatments? Cure, life-extension or symptom relief?

What is the success rate of these treatments?

When does treatment start?

Are there likely to be any side effects? How can I manage these?

Where will the treatments take place?

How long will the treatments last for?

Can I carry on life as usual during treatment?

Will the treatments affect my sex life?

Are there any clinical trials available?

## Questions to ask after treatments

---

How well did I respond to treatment?

What does the future hold?

What is the follow-up plan?

Are members of my family at higher risk for developing urothelial carcinoma because of my experience?

# Questions to ask about urothelial cancer

## Questions to ask about having surgery

What type of surgery will I have?

What will be removed?

How long will the surgery take?

How long will I be in the hospital?

How quickly will I be up and around again?

Does this surgery come with any long-term side effects?

How will the surgery affect my ability to urinate?

Will I need to have a urostomy? If so, how do I care for my urostomy?

If I get a neobladder, how will I know when to empty it?

## Questions to ask your insurer

Does my policy cover me for unfunded cancer treatment?

Does my policy cover me for my entire course of treatment? If not, which aspects of my treatment am I covered for?

How do I claim?

**KEYTRUDA (pembrolizumab) 50mg powder for infusion.** KEYTRUDA is a **Prescription Only Medicine Use:** KEYTRUDA KEYTRUDA is used in the treatment of melanoma which cannot be removed by surgery alone or when it has spread to multiple sites in the body in the treatment of a kind of lung cancer called non-small cell lung cancer (NSCLC). In the treatment of classical Hodgkin Lymphoma (cHL). In the treatment of urothelial carcinoma, including bladder cancer **Side effects:** Immune-mediated side effects including inflammation of the lungs, colon, liver, kidneys, pituitary gland, brain, eye, muscles, nervous system, pancreas, and heart, thyroid disorders, type 1 diabetes mellitus. Severe skin reactions including Steven-Johnson syndrome and toxic epidermal necrolysis. Severe infusion reactions including hypersensitivity and anaphylaxis. Transplant recipients: rejection of a transplanted organ, graft-versus-host-disease (in people with a bone marrow transplant using donor cells). Very common side effects include diarrhea, nausea, itching, rash, joint pain, back pain, feeling tired, cough, patches of discoloured skin, stomach pain, decreased levels of sodium in blood. Tiredness, nausea, vomiting, diarrhea, constipation, shortness of breath, rash, itching, headache, hair loss, and, infections of the upper respiratory tract were reported when given in combination with chemotherapy. You may experience more than one side effect at the same time. All medicines have risks and benefits. Talk to your doctor to see if KEYTRUDA is right for you. KEYTRUDA is a funded medicine for melanoma patients—restrictions apply. KEYTRUDA is an unfunded medicine for NSCLC, cHL and urothelial carcinoma patients. Ask your health professional the cost of the medicine and any other medical fees that may apply. Use only as directed and if symptoms continue or you have side effects, see your doctor, pharmacist, or health professional. Based on data sheet prepared 16 October 2017. Marketed by: Merck Sharp & Dohme (New Zealand) Limited, Newmarket, Auckland. For additional product information, consult the Consumer Medicine Information (CMI), available on request, phone 0800 500 673 or refer to the Medsafe website [www.medsafe.govt.nz](http://www.medsafe.govt.nz).

Copyright © 2017 Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Kenilworth, New Jersey, USA. All rights reserved. Copyright © 2017 Merck Sharp & Dohme (New Zealand) Limited. Level 3, 123 Carlton Gore Road, Newmarket, Auckland. All rights reserved. ONCO-1208503-0027 First Issued November 2017 TAPS NA9401 essence MSD8360